

TESTING PERTURBATIVE RESULTS WITH NON-PERTURBATIVE METHODS FOR THE HIERARCHICAL MODEL

Y. MEURICE and M. B. OKTAY *

*The University of Iowa, Department of Physics and Astronomy
Iowa City, Iowa 52242, USA*

We present non-perturbative methods to calculate accurately the renormalized quantities for Dyson's Hierarchical Model. We apply this method and calculate the critical exponent γ with 12 and 4 significant digits in the high and low temperature phases, respectively. We report accurate values for universal ratios of amplitudes and preliminary results concerning the comparison with perturbative results.

1. Introduction

The scalar field theory has many important applications such as the generation of mass in the standard model of elementary particles via spontaneous symmetry breaking and the theory of critical phenomena. However, there is no approximate treatment of the theory which can compete in accuracy with the perturbative treatment of quantum electrodynamics.

Hierarchical approximations, where the Renormalization Group (RG) transformations can be performed easily, provide a test ground for obtaining accurate calculations as well as for testing the validity of several perturbative techniques such as the renormalized perturbative expansion in field theory. Here, we present an approximation where simple algebraic methods can be used to calculate the renormalized quantities with great accuracy.

We consider Dyson's Hierarchical Model¹ that couples the main spin in boxes of size 2^l with a strength $(c/4)^l$ where c is a free parameter set to $c = 2^{1-2/D}$ in order to approximate a nearest neighbor model in D dimensions. We have to specify a local measure $W_0(\phi)$, for instance of the Landau-Ginsburg type $W_0(\phi) \propto e^{-[(1/2)m^2\phi^2 + \lambda_B\phi^4]}$ where the bare parameters will appear, or of the Ising type. Under a block spin transformation, the local measure changes according to

$$W_{n+1}(\phi) \propto e^{\frac{\beta}{2}(\frac{c}{4})^{n+1}} \times \int d\phi' W_n(\frac{\phi' - \phi}{2}) W_n(\frac{\phi' + \phi}{2}) \quad (1)$$

The recursion formula can be re-expressed in Fourier representation as

$$R_{n+1} \propto \exp\left(\frac{-1}{2}\beta\frac{\partial^2}{\partial k^2}\right)(R_n(\frac{\sqrt{c}k}{2}))^2 \quad (2)$$

*Presenting Author

where $R_n(k)$ is the Fourier transform of $W_n(\phi)$ with a rescaling of ϕ by a factor $2c^{-1/2}$.

2. Computational Method and Numerical Results

It was found that the finite dimensional truncations of degree l_{max} : $R_n(k) = 1 + a_{n,1}k^2 + a_{n,2}k^4 + \dots + a_{n,l_{max}}k^{2l_{max}}$ provide very accurate results in the symmetric phase^{2,3}. The round-off errors on the two point function grow like $(\beta - \beta_c)^{-1}$ and the finite truncations and volume effects can be controlled with an exponential precision. The non-Gaussian distribution of the numerical errors made in RG calculations related to small changes in the scaling factor was also studied in great detail⁴. The critical exponent associated with the two point function was found $\gamma = 1.299140730159 \mp 10^{-12}$. It was also shown that hyperscaling holds in $D = 3$ and that the coupling constant vanishes like $1/\Lambda$ in $D = 4$.

In the broken symmetry phase, it is possible to take advantage of about 10 iterations for which the low temperature scaling is observed⁵ to extrapolate to infinite volume limit. The critical exponents corresponding to zero momentum connected q -point functions $G_q^c(0)$, for $q = 1, 2, 3$ are $\gamma_1 = -0.324775 \mp 2 \times 10^{-5}$, $\gamma_2 = 1.29918 \mp 10^{-4}$ and $\gamma_3 = 2.928 \mp 10^{-2}$, respectively, in good agreement with the hyperscaling relation⁵

$$\gamma_q = \gamma[(q/2)\ln(4/c) - \ln 2]/\ln(2/c). \quad (3)$$

Recently, we have observed that the dimensionless renormalized coupling constants in the high temperature phase

$$\lambda_q^* = \frac{G_{2q}^c m_R^{q(2+D)-D}}{\beta^{(q-1)D/2}} \quad (4)$$

corresponding to the q -point function tend to universal values independent of the choice of the initial measure. Here G_{2q}^c is the connected Green function and m_R is the renormalized mass. We list below the values with significant digits common to an Ising and a Landau-Ginsburg measures.

Table 1. Renormalized coupling constants λ_q^* for q - point functions.

q	λ_q^*	q	λ_q^*
4	1.5058	8	579.97
6	18.107	10	35653

3. Perturbative Analysis

We will use the accurate results mentioned above to test perturbation theory. We expand the initial local measure into series as

$$e^{-(1/2)m^2\phi^2 - \lambda_B\phi^4} \simeq e^{-(1/2)m^2\phi^2} \left(1 - \lambda_B\phi^4 + \frac{\lambda_B^2}{2}\phi^8 + \dots\right). \quad (5)$$




Fig. 1. (a) Dimensionless Renormalized mass versus dimensionless bare mass; (b) Dimensionless Renormalized Coupling Constant versus dimensionless bare mass.

Substituting this back into the recursion relation, one obtains the recursion relation in orders of λ_B . After n iterations, the local measure becomes

$$W_n(\phi) = \exp\left[-\frac{(m_B^2 + k^2(n))}{2^{n+1}\phi^2}\right] \times (1 - \lambda_B \times \left(\frac{3}{2^n}\phi^2 \sum_{m=0}^{n-1} \frac{2^{-m}}{(m_B^2 + k^2(m))} + \frac{\phi^4}{2^{3n}}\right) + \vartheta(\lambda_B^2)) \quad (6)$$

where m_B^2 stands for the bare mass and $k^2(l)$ can be interpreted as the momentum square over l -iterations. We have calculated the renormalized mass and the dimensionless coupling constant up to order of λ_B^3 . The details of the results will be provided soon⁶. This is illustrated in the figures above. The iterative method can be used to high order in λ_B if one does not insist on obtaining closed form as in Eq. (6). This will be used to test standard perturbation method⁷ to evaluate the critical exponents.

References

1. F. Dyson, *Comm. Math. Phys* **12**, 91 (1969) ; G. Baker, *Phys. Rev.* **B5**, 2622 (1972).
2. J. J. Godina, Y. Meurice, S. Niermann and M. B. Oktay, *Phys. Rev.* **D57**, 6326 (1998).
3. J. J. Godina, Y. Meurice and M. B. Oktay, *Phys. Rev.* **D59**, 096002 (1999).
4. Y. Meurice and M. B. Oktay, hep-lat/0005011, *Phys. Rev.* **D** in press.
5. J. J. Godina, Y. Meurice and M. B. Oktay, *Phys. Rev.* **D61**, 114509 (2000).
6. Y. Meurice and M. B. Oktay, U. Of Iowa preprints in preparation.
7. G. Parisi, *Statistical Field Theory* (Addison-Wesley, New York, 1988).